



UPPSALA
UNIVERSITET

Chronic stress and the adult zebrafish brain: the importance of the GABA signaling system by a gene-expression perspective

1. Introduction

Animals live in an environment that poses fundamental surviving challenges, which need to be sensed and interpreted to produce proper behavioral responses. In the vertebrate brain, potentially life-threatening conditions cause the activation of defensive survival circuits.¹ Those involve descending crosstalks between the cortex and the amygdala, and the latter and the hypothalamus, as well as ascending projections traveling the opposite direction.²⁻⁷ Descending brainstem projections affect the activity of hypothalamic nuclei, too.⁴⁻⁶ The ultimate effect of the activation of defensive survival circuits is a switch-on of the endocrine stress response, achieved through communications between the hypothalamus and the hypophysis.^{1,4,5} This starts with the release of corticotropin-releasing hormone (CRH) by CRH cells located in the paraventricular nucleus (PVN) of the hypothalamus. CRH in turn stimulates the secretion of adrenocorticotrophic hormone (ACTH) from the anterior pituitary, which subsequently stimulates the synthesis and systemic secretion of glucocorticoids from the adrenal cortex into the bloodstream^{4,5}. Glucocorticoids (mainly cortisol) regulate the energy status of the body, promoting energy mobilization and therefore facilitating a reaction to a threat.⁶ The physiological pathway, starting from the nervous system and eventually affecting the adrenal cortex, is termed hypothalamo-pituitary-adrenal (HPA) axis in mammals.⁶ In teleosts, no condensed adrenal gland is present; it is rather possible to identify sparse cell clusters analogous either to the adrenal medulla or cortex.⁸ Thus, for fish the definition hypothalamo-pituitary-interrenal (HPI) axis is used.

In the adult vertebrate brain, inhibition of defensive survival circuits is mediated by the major inhibitory neurotransmitter γ -aminobutyric acid (GABA), which is synthesized from α , L-glutamate by glutamate decarboxylases (GADs).^{9,10} GABA is widely employed as signaling molecule in defensive survival circuits, as reported by immunological studies on GADs distribution in both mammals and fish.¹⁰⁻¹³ The use of GABA is also witnessed by GABA_A receptor subunits, which assemble in a region-specific fashion with a strict stoichiometry.¹⁴⁻¹⁹ GABA_A receptors are homo- or heteropentamers; in mammals there are nineteen genes encoding for its monomers, in zebrafish at least twenty-three.^{20,21} In the amygdala of rats and mice α_2 and α_3 are detected to the highest level among the α monomers.^{14,17-19} The β_3 and γ_1 subunits are also produced to high levels; in this brain area $\alpha_2\beta_3\gamma_1$, $\alpha_3\beta_3\gamma_1$ GABA_A receptors are found. The γ_2 monomer is present as well, and receptors incorporating this subunits may also be located in the amygdala.^{14,19} In the hippocampus $\alpha_5\beta_3\gamma_2$ GABA_A channels are present.^{14,15,19,22} The pattern of GABA_A receptor subunits expression is somewhat different between rats and mice in the hypothalamus. In the former species the $\alpha_2\beta_3\gamma_1$ combination is detected, whereas the latter also presents the α_2 , β_1 , and γ_2 monomers detected to a certain level.^{14,19} Chronic stress impairs GABA signaling in

the defensive survival circuits, both at the gene expression level^{23,24} and in terms of altered chloride permeability of GABA target cells.²⁵

Also the HPA-axis is under GABAergic control. CRH neurons in the PVN receive tonic inhibitory GABAergic input that constrains CRH neuronal activity under basal conditions.²⁶ Following an acute threat, the frequency of spontaneous inhibitory postsynaptic currents is reduced up to 5 h later, causing an increase in circulating glucocorticoid levels. Chronic stress and sustained increases in systemic glucocorticoid levels lead to suppression of GABAergic inhibition of CRH neurons, by a reduction in GABA synapses.²⁶ Also the composition of the GABA_A receptors in the PVN can change in response to chronic stress, towards increased expression of neurosteroid-sensitive GABA_A receptor δ subunit-containing receptors.²⁴ Stress-derived steroid hormones can be metabolized to neurosteroids (such as THDOC and allopregnanolone), which can act as positive allosteric modulators of GABA_A receptor δ subunit-containing receptors, potentiating the tonic component of GABAergic inhibition.²⁷ This would provide a reciprocal regulation of stress hormones and GABA receptors, where GABAergic inhibition regulates the HPA-axis and the production of stress hormones and derived neurosteroids also alter this GABAergic inhibition during chronic stress.²⁷

2. Aim of the study in detail

The aim of the project is to investigate the effect of acute versus chronic stress on the GABA signaling system in the adult zebrafish brain. Our group has recently characterized the GABA signaling system in the adult zebrafish brain²¹, and qPCR primers have been developed to quantify mRNA expression of GABA_A receptor subunits and GADs in specific brain areas. Both female and male zebrafish will be studied, allowing for investigation of potential sex differences.

3. Techniques

- a. Dissection of the zebrafish brain;
- b. RNA extraction and RNA quality evaluation;
- c. Reverse transcription of RNA into cDNA;
- d. RT-qPCR;
- e. Measurement of water total cortisol;
- f. Measurement of plasma cortisol;
- g. Data analysis.

4. Supervisors

Prof. Svante Winberg (Uppsala University, Sweden) <svante.winberg@neuro.uu.se>, Dr. Arianna Cocco (formerly Uppsala University, Sweden, now Institute of Science and Technology Austria), Dr Per-Ove Thörnqvist (Uppsala University, Sweden).

5. References

1. LeDoux JE (2014) Coming to terms with fear. *Proc Natl Acad Sci U S A* **111**(8):2871-2878
2. Papez JW (1937). A proposed mechanism of emotion. *Arch Neurol Psychiat* **38**:725-743
3. Sah P, Faber SL, Lopez de Armentia M, Power J (2003) The Amygdaloid Complex: Anatomy and Physiology. *Physiol Rev* **83**(3):803-834

4. Gray H (2005) Chapter 21: Diencephalon. In: Gray's Anatomy: The Anatomical Basis of Clinical Practice, 39th edition, AR Crossman and S Standring (eds.), pgg. 369-385, Elsevier Churchill Livingstone, © Elsevier Ltd
5. Herman JP, Ostrander MM, Mueller NK, Figueiredo H (2005) Limbic system mechanisms of stress regulation: Hypothalamo-pituitary-adrenocortical axis. *Prog Neuropsychopharmacol Biol Psychiatry* **29**(8):1201-1213
6. Herman JP, McKleeven JM, Ghosal S, Kopp B, Wulsin A, Makinson R, Scheimann J, Myers B (2016) Regulation of the hypothalamic-pituitary-adrenocortical stress response. *Compr Physiol* **6**(2):603-621
7. LeDoux JE, Brown R (2017) A higher order theory of emotional consciousness. *Proc Natl Acad Sci U S A* **114**(10):E2016-E2025
8. Clark KJ, Boczek NJ, Ekker SC (2011) Stressing zebrafish for behavioral genetics. *Rev Neurosci* **22**(1):49-62
9. Robert E, Kuriyama K (1968) Biochemical-physiological correlations in studies of the γ -aminobutyric acid system. *Brain Res* **8**(1):1-35
10. Tappaz ML, Brownstein MJ, Kopin IJ (1977) Glutamate decarboxylase (GAD) and γ -aminobutyric acid (GABA) in discrete nuclei of hypothalamus and substantia nigra. *Brain Res* **125**(1):109-121
11. Sheikh SN, Martin SB, Martin DL (1999) Regional distribution and relative amounts of glutamate decarboxylase isoforms in rat and mouse brain. *Neurochem Int* **35**(1):73-80
12. Anglade I, Mazurias D, Douard V, Le Jossic-Corcus C, Mañanos EL, Michel D, Kah O (1999) Distribution of Glutamic Acid Decarboxylase mRNA in the Forebrain of the Rainbow Trout as Studied by In Situ Hybridization. *J Comp Neurol* **410**(2):277-289
13. Martyniuk CJ, Awad R, Hurley R, Finger TE, Trudeau VL (2007) Glutamic acid decarboxylase 65, 67, and GABA-transaminase mRNA expression and total enzyme activity in the goldfish (*Carassius auratus*) brain. *Brain Res* **1147**:154-166
14. Wisden W, Laurie DJ, Monyer H, Seeburg PH (1992) The Distribution of 13 GABA_A Receptor Subunit mRNAs in the Rat Brain. I. Telencephalon, Diencephalon, Mesencephalon. *J Neurosci* **12**(3):1040-1062
15. Somogyi P, Fritschy J-M, Benke D, Roberts JDB, Sieghart W (1996) The γ 2 Subunit of the GABA_A Receptor is Concentrated in Synaptic Junctions Containing the α 1 and β 2/3 Subunits in Hippocampus, Cerebellum and Globus Pallidus. *Neuropharmacology* **35**(9-10):1425-1444
16. Wei W, Zhang N, Peng Z, Houser CR, Mody I (2003) Perisynaptic Localization of δ Subunit-Containing GABA_A Receptors and Their Activation by GABA Spillover in the Mouse Dentate Gyrus. *J Neurosci* **23**(33):10650-10661
17. Marowsky A, Fritschy J-M, Vogt KE (2004) Functional mapping of GABA_A receptor subtypes in the amygdala. *Eur J Neurosci* **20**(5):1281-1289
18. Marowsky A, Rudolph U, Fritschy J-M, Arand M (2012) Tonic Inhibition in Principal Cells of the Amygdala: A Central Role for α 3 Subunit-Containing GABA_A Receptors. *J Neurosci* **32**(25):8611-8619
19. Hörtnagl H, Tasan RO, Wieselthaler A, Kirchmair E, Sieghart W, Sperk G (2013) Patterns of mRNA and protein expression for 12 GABA_A receptor subunits in the mouse brain. *Neuroscience* **236**:345-372
20. Simon J, Wakimoto H, Fujita N, Lalande M, Barnard EA (2004) Analysis of the Set of GABA_A Receptor Genes in the Human Genome. *J Biol Chem* **279**(40):41422-41435
21. Cocco A, Rönnberg AMC, Jin Z, André IG, Vossen LE, Bhandage AK, Thörnqvist P-O, Birnir B, Winberg S (2017) Characterization of the γ -aminobutyric acid signaling system in the zebrafish (*Danio rerio* Hamilton) central nervous system by reverse transcription-quantitative polymerase chain reaction. *Neuroscience* **343**:300-321
22. Caraiscos VB, ElliotEM, You-Ten KE, Cheng VY, Beelli D, Newell JG, Jackson MF, Lambert JJ, Roshal TW, Wafford KA, MacDonald JF, Orser BA (2004) Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by α 5 subunit-containing γ -aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A* **101**(10):3662-3667
23. Cullinan WE, Wolfe TJ (2000) Chronic stress regulates levels of mRNA transcripts encoding β subunits of the GABA_A receptor in the rat stress axis. *Brain Res* **887**:118-124
24. Verkuyl JM, Hemby SE, Joël M (2004) Chronic stress attenuates GABAergic inhibition and alters gene expression of parvocellular neurons in rat hypothalamus. *Eur J Neurosci* **20**(6):1665-1673
25. Hewitt SA, Wamsteeker JI, Kurz EU, Bains JS (2009) Altered chloride homeostasis remove synaptic inhibitory constraint of the stress axis. *Nat Neurosci* **12**(4):438-443
26. Levy BH, Tasker JG (2012) Synaptic regulation of the hypothalamic-pituitary-adrenal axis and its modulation by glucocorticoids and stress. *Front Neurosci* **6**: 1-13

27. Mody I, Maguire J (2012) The reciprocal regulation of stress hormones and GABA_A receptors.
Front Cell Neurosci **6**: 1-6